

STEENROD SQUARES ON CONJUGATION SPACES

MATTHIAS FRANZ AND VOLKER PUPPE

ABSTRACT. We prove that the coefficients of the so-called conjugation equation for conjugation spaces in the sense of Hausmann–Holm–Puppe are completely determined by Steenrod squares. This generalises a result of V. A. Krasnov for certain complex algebraic varieties. It also leads to a generalisation of a formula given by Borel and Haefliger, thereby largely answering an old question of theirs in the affirmative.

1. STATEMENT OF THE RESULTS

Let X be a topological space with an involution τ . We look at X as a space with an action of the group $G = \{1, \tau\}$. We take cohomology with coefficients in $\mathbf{F}_2$ and consider the restriction map $r: H^*(X) \rightarrow H^*(X^\tau)$, its equivariant counterpart $r_G: H_G^*(X) \rightarrow H_G^*(X^\tau) = H^*(X^\tau) \otimes \mathbf{F}_2[u]$ and the canonical projection $p: H_G^*(X) \rightarrow H^*(X)$.

According to Hausmann–Holm–Puppe [HHP], (X, τ) is called a *conjugation space* if there exists a section $\sigma: H^*(X) \rightarrow H_G^*(X)$ to p and a degree-halving isomorphism $\kappa: H^*(X) \rightarrow H^*(X^\tau)$ with the following property: for every $x \in H^n(X)$, $n \in \mathbf{N}$, there exists elements $y_1, \dots, y_n \in H^*(X^\tau)$ such that the so-called *conjugation equation*

$$(1) \quad r_G(\sigma(x)) = \kappa(x)u^n + y_1u^{n-1} + \dots + y_{n-1}u + y_n$$

holds. A priori, σ and κ are only assumed to be additive, but the conjugation equation implies that they are in fact multiplicative and unique.

The coefficients y_i in the conjugation equation are completely determined by x . We prove that they are the Steenrod squares of $\kappa(x)$.

Theorem 1.1. *For every $x \in H^{2n}(X)$, $n \in \mathbf{N}$, one has*

$$r_G(\sigma(x)) = \sum_{i=0}^n \text{Sq}^i(\kappa(x)) u^{n-i}.$$

Corollary 1.2. *For every $x \in H^*(X)$, one has*

$$r(x) = \kappa(x)^2.$$

We also show that the isomorphism κ commutes with total Steenrod squares.

Theorem 1.3. *For every $x \in H^*(X)$ one has*

$$\kappa(\text{Sq}(x)) = \text{Sq}(\kappa(x)).$$

Note that the odd Steenrod squares of x vanish since $H^*(X)$ is concentrated in even degrees. Hence, the above identity is equivalent to

$$(2) \quad \kappa(\text{Sq}^{2k}(x)) = \text{Sq}^k(\kappa(x)) \quad \text{for all } k \in \mathbf{N}.$$

2. PROOFS

We denote the Steenrod algebra for the prime 2 by $\mathcal{A}$.

Lemma 2.1. *For every n there exist elements $a_0, \dots, a_n, b \in \mathcal{A}$ such that for every conjugation space X and every $x \in H^{2n}(X)$ one has*

$$r_G(\sigma(x)) = \sum_{i=0}^n a_i(\kappa(x)) u^{n-i} \quad \text{and} \quad \kappa(\text{Sq}(x)) = b(\kappa(x)).$$

Moreover, $a_0 = 1$ and $a_1 = \text{Sq}^1$.

Proof. Since κ is bijective, one can define functions a_i and b for each X such that the above identities hold. We show that they are (or, more precisely, come from) Steenrod squares, using that the restriction map r_G commutes with all Steenrod squares. We write $\kappa(x) = z$.

We start by proving the claim about the a_i by induction, beginning at $a_0(z) = z$. If $i > 0$ is even, we apply Sq^{2k} , where $k \leq i/2$ will be chosen later. We can write

$$(3) \quad \text{Sq}^{2k}(\sigma(x)) = \sum_{l=-n}^k \sigma(\tilde{x}_l) u^{2(k-l)}$$

for some $\tilde{x}_l \in H^{2(n+l)}(X)$. Write $z_l = \kappa(\tilde{x}_l)$. The restriction $r_G(\sigma(\tilde{x}_l) u^{2(k-l)})$ has leading term $z_l u^{n+2k-l}$, while the leading power of u in $\text{Sq}^{2k}(r_G(\sigma(x)))$ is at most u^{n+2k} . Hence, the summation in (3) is in fact only over $0 \leq l \leq k$.

Comparing coefficients of u^{n+2k-l} yields

$$(4) \quad z_l = \sum_{j=0}^l \binom{n-l+j}{2k-j} \text{Sq}^j(a_{l-j}(z)) + \sum_{j=1}^l \text{Sq}^j(z_{l-j}).$$

This shows $z_l = b_l(z)$ for some $b_l \in \mathcal{A}$. Comparing coefficients of u^{n+2k-i} gives

$$\begin{aligned} \sum_{l=0}^k a_{i-l}(z_l) &= \binom{n}{2k} a_i(z) + \sum_{l=1}^k a_{i-l}(b_l(z)) \\ &= \binom{n-i}{2k} a_i(z) + \sum_{j=1}^{2k} \binom{n-i+j}{2k-j} \text{Sq}^j(a_{i-j}(z)). \end{aligned}$$

Now suppose that $k \leq i/2$ is such that

$$\binom{n}{2k} \neq \binom{n-i}{2k}.$$

For instance, this is true if $2k$ is the largest power of 2 dividing i . (Recall that a binomial coefficient mod 2 is the product of the binomial coefficients taken for each pair of binary digits, cf. [S, Lemma I.2.6].) Then the above equation can be solved for $a_i(z)$ and shows that $a_i(z)$ can be expressed in terms of repeated Steenrod squares of z .

For odd i , a similar (but simpler) reasoning based on commutativity with respect to Sq^1 gives $a_i(z) = \text{Sq}^1(a_{i-1}(z))$, in particular $a_1(z) = \text{Sq}^1(z)$.

Now that all $a_i(z)$ are known, we apply Sq^{2k} for any k . Using the same notation as above, we have $\text{Sq}^{2k}(x) = p(\text{Sq}^{2k}(\sigma(x))) = \tilde{x}_k$. Comparing coefficients as before gives a formula for $b_l(z)$ similar to equation (4), but where the summation index j starts at $l-n$ if $l > n$. Still, the equations can be recursively solved for z_l . Hence,

$$\kappa(\text{Sq}(x)) = \kappa(\tilde{x}_0) + \dots + \kappa(\tilde{x}_n) = b_0(z) + \dots + b_n(z) = b(z). \quad \square$$

In principle, the preceding proof could be used to determine the coefficients of the conjugation equation completely (as well as those of $\text{Sq}^k(\sigma(x))$ for any k). We will take a less tedious approach which relies on the fact that suitable products

of infinite-dimensional real projective space can “detect” Steenrod squares, cf. [S, Corollary I.3.3].

Fact 2.2. *The restricted evaluation map*

$$\mathcal{A}_{\leq n} \rightarrow H^*((\mathbf{RP}^\infty)^n), \quad a \mapsto a(u \times \cdots \times u)$$

is injective for any $n \in \mathbf{N}$.

Proof of Theorem 1.1. Write SQ for the homogenised total Steenrod square,

$$\text{SQ}(\kappa(x)) = \sum_{i=0}^n \text{Sq}^i(\kappa(x)) u^{n-i}.$$

We want to show $r_G(\sigma(x)) = \text{SQ}(\kappa(x))$ for all cohomology classes of all conjugation spaces. By Lemma 2.1 and Fact 2.2, it suffices to do so for the conjugation space $X = (\mathbf{CP}^\infty)^n$ with complex conjugation and x , the n -fold cross product of the generator $v \in H^2(\mathbf{CP}^\infty)$ because $X^\tau = (\mathbf{RP}^\infty)^n$ and $\kappa(x) = u \times \cdots \times u$ in this case. For $n = 1$ the identity is true since we already know a_1 . The general case reduces to the case $n = 1$ because of the multiplicativity of all maps involved:

$$\begin{aligned} r_G(\sigma(x)) &= r_G(\sigma(v \times \cdots \times v)) = r_G(\sigma(v)) \times \cdots \times r_G(\sigma(v)) \\ &= \text{SQ}(\kappa(v)) \times \cdots \times \text{SQ}(\kappa(v)) = \text{SQ}(\kappa(v \times \cdots \times v)) = \text{SQ}(\kappa(x)). \end{aligned}$$

(Note that κ , σ and r_G commute with products.) □

Proof of Corollary 1.2. We have for $x \in H^{2n}(X)$

$$r(x) = r(p(\sigma(x))) = p(r_G(\sigma(x))) = p(\text{Sq}^n(\kappa(x))) = \kappa(x)^2. \quad \square$$

Proof of Theorem 1.3. As in the proof of Theorem 1.1, it suffices to show the claimed identity for $X = (\mathbf{CP}^\infty)^n$ and $x = v \times \cdots \times v$. Again, the general case can be reduced to $n = 1$, where we find

$$\kappa(\text{Sq}(v)) = \kappa(v + v^2) = \kappa(v) + \kappa(v)^2 = \text{Sq}(\kappa(v)). \quad \square$$

3. REMARKS

Let X be a non-singular complex projective variety defined over the reals such that its real locus X^τ is non-empty. All algebraic cycles in X are understood to be defined over the reals. Borel–Haefliger have shown that if $H_*(X)$ and $H_*(X^\tau)$ are generated by algebraic cycles, then the restriction λ of cycles in X to their real locus induces a degree-halving isomorphism $H_*(X) \rightarrow H_*(X^\tau)$ respecting intersection products [BH, §5.15]. They also show that if $H^*(X^\tau)$ is generated by algebraic cycles and $x \in H^*(X)$ is Poincaré dual to a linear combination of non-singular subvarieties, then the identity in Theorem 1.3 holds, and they ask whether it holds more generally [BH, §5.17].

V. A. Krasnov has proved that for a variety X as above, Theorem 1.1 holds for cohomology classes Poincaré dual to algebraic cycles, where κ is the Poincaré transpose of λ and σ the canonical section [K1, Theorem 4.2]. This implies that if $H_*(X)$ is generated by algebraic cycles, then so is $H_*(X^\tau)$ [K2, Theorem 0.1]. Moreover, X is a conjugation space in the sense of [HHP].

In a topological framework van Hamel has recently shown that certain topological manifolds with involutions are conjugation spaces [vH, Theorem]. The necessary assumptions are formulated in terms of topological cycles.

The following simple example shows that in general the existence of a degree-halving multiplicative isomorphism $\kappa: H^*(X) \rightarrow H^*(X^\tau)$ by itself does not imply that (X, τ) is a conjugation space.

Example 3.1. Let $X = S^2 \times S^4$ be equipped with the componentwise involution τ which is the identity for S^2 and for S^4 has fixed point set S^1 . So $X^\tau = S^2 \times S^1$. Clearly there is a degree-halving multiplicative isomorphism $\kappa: H^*(X) \rightarrow H^*(X^\tau)$. It is easy to check there is also a multiplicative section $\sigma: H^*(X) \rightarrow H_G^*(X)$. But (X, τ) is not a conjugation space: the restriction map

$$r_G: H_G^*(S^2 \times S^4) \cong H^*(S^2 \times S^4) \otimes \mathbf{F}_2[u] \rightarrow H^*(S^2 \times S^1) \otimes \mathbf{F}_2[u]$$

is given by $s_2 \otimes 1 \mapsto s_2 \otimes 1$ and $s_4 \otimes 1 \mapsto s_1 \otimes u^3$, where $s_n \in H^n(S^n)$ denotes the generator. Hence the conjugation equation does not hold. Of course, $S^2 \times S^4$ with the different componentwise involution $\tilde{\tau}$ which has $S^1 \subset S^2$ and $S^2 \subset S^4$ as fixed point sets (and hence $X^{\tilde{\tau}} = S^1 \times S^2 \cong X^\tau$) is a conjugation space.

REFERENCES

- [BH] A. Borel and A. Haefliger, La classe d'homologie fondamentale d'un espace analytique, Bull. Soc. Math. France **89** (1961), 461–513
- [HHP] J.-C. Hausmann, T. Holm, V. Puppe, Conjugation spaces, Alg. Geom. Topology **5** (2005), 923–964
- [K1] V. A. Krasnov, On equivariant Grothendieck cohomology of a real algebraic variety, and its applications, Russian Acad. Sci. Izv. Math. **44** (1994), 461–477
- [K2] V. A. Krasnov, Real algebraically maximal varieties, Math. Notes **73** (2003), 806–812
- [S] N. E. Steenrod, Cohomology operations, Princeton University Press, Princeton, N. J. 1962
- [vH] J. van Hamel, Geometric cohomology frames on Hausmann–Holm–Puppe conjugation spaces, ArXiv e-print math.AT/0509498 (2005)

E-mail address: `matthias.franz@ujf-grenoble.fr`

E-mail address: `volker.puppe@uni-konstanz.de`

FACHBEREICH MATHEMATIK, UNIVERSITÄT KONSTANZ, 78457 KONSTANZ, GERMANY